

High-Spin Octupole Correlations in the $N = 85$, ^{139}Xe and ^{141}Ba Isotones

*S.J. Zhu, J.H. Hamilton, A.V. Ramayya, M.G. Wang,
Physics Department, Vanderbilt University, Nashville, TN 37235
J.O. Rasmussen, I.Y. Lee, S.Y. Chu
Lawrence Berkeley National Laboratory, Berkeley, CA 94720
M.A. Stoyer
Lawrence Livermore National Laboratory, Livermore, CA 94550
Others of the GANDS95 Collaboration*

Prompt γ -rays from the fission fragments of ^{252}Cf spontaneous fission were investigated. Data presented in this work from the 1995 Gammasphere measurements of the GANDS95 collaboration [1] were analyzed mainly by Vanderbilt University members of the collaboration. The results of this work were presented in ref. [2].

High-spin states in ^{139}Xe and ^{141}Ba were investigated using gamma-gamma and gamma-gamma-gamma coincidence studies of spontaneous fission of ^{252}Cf . Theoretical calculations [3] and comparison of the odd-even staggering of the differential radii [4] indicate that octupole correlations may occur in ^{139}Xe and ^{141}Ba . On the other hand, some heavier $N = 85$ isotones like ^{145}Nd and ^{147}Sm , have been described as ‘quasi- $f_{7/2}$ ’ nuclei. Octupole and ‘quasi- $f_{7/2}$ ’ bands are observed in these nuclei. Of particular interest is a new band assigned to have positive parity and based on a $\nu i_{13/2}(f_{7/2})^2$ multiplet in both isotones. The systematics in the $N = 85$ isotones for these ‘quasi- $f_{7/2}$ ’ states are presented in Figure 1. Alternating parity bands built on a $7/2^-$ level are assigned up to spin $37/2$ and $29/2$ in ^{139}Xe and ^{141}Xe , respectively. Another possible octupole band with spin up to $29/2$ also is observed in both nuclei. Cascades of enhanced $E1$ transitions and $B(E1)/B(E2)$ ratios indicate strong octupole correlations.

References

- [1] For GANDS95 list of authors and institutions see B.R.S. Babu, *et al.*, Phys. Rev. **C54** (1996) 568.
- [2] S.J. Zhu, *et al.*, J. of Phys. **G23** (1997) L77.
- [3] G.A. Leander, *et al.*, Phys. Lett. **152B** (1985) 284, and S. Cwiok and W. Nazarewicz, Nucl. Phys. **A469** (1989) 367.
- [4] R. Sheline, K. Jian and K. Jian, Phys. Rev. **C38** (1988) 2925.




Fig. 1. Level systematics for selected low-lying levels in $N = 85$ isotopes. The level energies are given relative to the first $7/2^+$ state.